



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 263/08	A1	(11) International Publication Number: WO 90/02738 (43) International Publication Date: 22 March 1990 (22.03.90)
(21) International Application Number: PCT/US89/03827 (22) International Filing Date: 12 September 1989 (12.09.89) (30) Priority data: 244,126 14 September 1988 (14.09.88) US (60) Parent Application or Grant (63) Related by Continuation US 244,126 (CIP) Filed on 14 September 1988 (14.09.88) (71) Applicant (for all designated States except US): SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033 (US). (72) Inventors; and (75) Inventors/Applicants (for US only) : SCHUMACHER, Doris, P. [US/US]; 76 Edgewood Drive, Florham Park, NJ 07932 (US). CLARK, Jon, E. [US/US]; 501 Grove Avenue, Highland Park, NJ 08904 (US). MURPHY, Bruce, L. [US/US];		969 Bloomfield Avenue, Apartment A7, Glen Ridge, NJ 07028 (US). (74) Agents: MAJKA, Joseph, T. et al.; Schering-Plough Corporation, One Giralda Farms, Madison, NJ 07940-1000 (US). (81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CF (OAPI patent), CG (OAPI patent), CH (European patent), CM (OAPI patent), DE (European patent), DK, FI, FR (European patent), GA (OAPI patent), GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL (European patent), NO, RO, SD, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent), US. Published <i>With international search report.</i>
(54) Title: PROCESS FOR PREPARING OXAZOLINE COMPOUNDS (57) Abstract A novel process for preparing oxazoline compounds is disclosed. The process utilizes an amino alcohol, a cyano compound, a base and a dihydric alcohol solvent, a polyhydric alcohol solvent or mixtures thereof.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Fasso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

PROCESS FOR PREPARING OXAZOLINE COMPOUNDS

FIELD OF THE INVENTION

The present invention is directed to a novel process for preparing oxazoline compounds. Oxazoline compounds are useful in preparing surface active agents, detergents, waxes, and intermediates for pharmaceutical compounds, such as those disclosed in U.S. Patent 4,743,700.

BACKGROUND OF THE INVENTION

Oxazoline compounds are known, and are disclosed in U.S. Patents 4,216,162; 3,813,378; 2,786,870; 2,718,520; and 2,820,041.


Processes for making oxazolines are known, such as described in European Patent Application 130,633, U.S. Patent 4,235,892 and the article "Formation of Cyclic Imidic Esters by Reaction of Nitriles with Amino Alcohols" by H. Witte and Wolfgang Seeliger, Liebigs Ann Chem. pp 996-1009, (1974) which utilizes catalytic amounts of certain metal salts. U.S. Patent 2,402,198 describes a process for preparing oxazolines by reacting monoethanolamine with a nitrile in the presence of an alkaline catalyst. U.S. Patent 2,759,001 discloses preparing racemic mixtures of isomeric oxazolines by

-2-

reacting dichloroacetonitrile with an aminodiol compound. U.S. Patent 3,979,405 discloses preparing 2-oxazolines by reacting an amino alcohol with a nitrile in an anhydrous alcohol such as n-butanol or cyclohexanol. None of these references teach a method for preparing oxazolines employing a dihydric or polyhydric alcohol solvent. It would be desirable to provide a process for preparing oxazoline compounds whose yields, purity and selectivity are as good as or better than methods previously taught. It would also be desirable to provide a process for preparing said oxazoline compounds which requires as few or even fewer steps than methods previously taught. It would also be desirable to provide a process which is as economical, if not more so, than previous methods. Further, where two or more potential structural isomers may be formed, it would be highly desirable to provide a process which is regio-selective i.e. produces only one structural isomer.

SUMMARY OF THE INVENTION

The present invention is directed toward a process for preparing oxazoline compounds of the formula:



wherein R^1 , R^2 , R^3 , R^4 and R^5 can be the same or different, each independently represents


-3-

hydrogen, alkyl, haloalkyl, cycloalkyl or substituted cycloalkyl, cycloalkalkyl or substituted cycloalkalkyl, alkenyl or substituted alkenyl, alkynyl, alkenylalkyl or substituted alkenylalkyl, alkynylalkyl, alkoxyalkyl or substituted alkoxyalkyl, dialkylamino-alkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, arylalkenyl or substituted arylalkenyl, alkoxyaryl or substituted alkoxyaryl, aryloxyaryl or substituted aryloxyaryl, aryloxyalkyl or substituted aryloxyalkyl, acyl or substituted acyl, aromatic heterocyclic or substituted aromatic heterocyclic, heterocyclic alkyl or substituted heterocyclic alkyl, heterocyclic cycloalkyl or substituted heterocyclic cycloalkyl, heterocyclic cycloalkylalkyl or substituted heterocyclic cycloalkylalkyl, sulfoxide or substituted sulfoxide, sulfonyl or substituted sulfonyl, sulfide or substituted sulfide or hydroxyalkyl.

The process comprises the step of contacting a cyano compound of the formula



with an aminoalcohol compound of the formula




or salt thereof, wherein R^1 , R^2 , R^3 , R^4 and R^5 are as defined hereinbefore, performed in the presence of a base and a dihydric alcohol solvent, a polyhydric alcohol solvent or mixture thereof. Preferably R^1 and R^2 independently represent hydrogen, sulfide, sulfoxide, sulfonyl, aryl or substituted aryl; R^3 and R^4

-4-

independently represent hydrogen, alkyl or hydroxyalkyl; and R⁵ is phenyl, 4-nitrophenyl, cinnamyl, 4-methoxycinnamyl or dichloromethyl. More preferably, R² is 4-methylthiophenyl, 4-methyl-SO-phenyl or 4-methyl-SO₂-phenyl; R⁴ is hydroxymethyl and R⁵ is phenyl or dichloromethyl when R¹ and R³ are hydrogen. Preferably the base is diazabicycloundecene, more specifically 1,8-diazobicyclo[5.4.0]undec-7-ene (DBU) or an alkali metal carbonate, most preferably potassium carbonate. Also, preferred is that the alcohol solvent is a mixture of a dihydric alcohol and a polyhydric alcohol, most preferably a mixture of ethylene glycol and glycerol.

Also preferred are those amino alcohol starting materials, (IX) of the general formula



wherein R¹ and R² are as defined hereinabove. For these starting materials the process is regio-selective. That is, although the reaction potentially can give rise to two or more structural isomers, the present process produces only one of the two or more potential structural isomers.

The process has the advantages of being able to prepare an oxazoline compound of formula (X) in high yields, good purity, high specificity, with low by-product formation using relatively mild reaction conditions with as few or fewer steps than other processes previously taught. The present invention has the further advantage of providing a process of preparing oxazoline compounds as economically, if not more so, than other processes previously taught. The present invention has the further advantage of providing a process for

-5-

preparing oxazoline compounds whose stereoisomeric configuration can be easily determined beforehand simply by the selection of the appropriate starting materials. An especially advantageous feature of the present invention is that virtually no racemization occurs when chiral starting material of formula IX are used. In the particular situation where the amino alcohol starting material contains two hydroxyl groups, the present process has the advantage in that it is regio-selective.

DETAILED DESCRIPTION OF THE EMBODIMENTS

When utilized in the present specification and in the appended claims the terms listed hereinbelow, unless otherwise indicated are defined as follows:

The term "alkyl" refers to a straight saturated hydrocarbon moiety (i.e. hydrocarbons having carbon-carbon single bonds) containing from 1 to 6 carbon atoms, or a branched saturated hydrocarbon moiety of 3 to 6 carbon atoms, such as for example, methyl (i.e. $-\text{CH}_3$), ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl and the like; the term "substituted alkyl" refer to an alkyl moiety which if further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano (i.e. $-\text{CN}$), carboxyl (i.e. $-\text{COOH}$) or salts thereof, nitro (i.e. $-\text{NO}_2$) and hydroxyl;

The terms "halogen" and "halo" refers to fluoride, chloride, bromide or iodide.

The term "haloalkyl" refers to an alkyl moiety in which one or more of the hydrogen atoms has been replaced by a halogen atom, such as, for example, chloromethyl, fluoromethyl, bromomethyl, trifluoromethyl, dichloromethyl, 2-chloro-2-fluoroethyl, 6,6,6-trichlorohexyl and the like.

-6-

The term "cycloalkyl" refers to a saturated carbocyclic ring characterized by closed rings and containing from 3 to 6 carbon atoms, such as for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like; the term "substituted cycloalkyl" refers to a cycloalkyl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl, or salts thereof, nitro and hydroxyl;

The term "cycloalkalkyl" refers to a cycloalkyl moiety of 3 to 6 carbon atoms covalently bonded to an alkyl moiety of 1 to 6 carbon atoms; the term substituted "cycloalkalkyl" refer to a cycloalkalkyl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano (i.e. -CN), carboxyl (i.e. -COOH) or salts thereof, nitro (i.e. -NO₂) and hydroxyl;

The term "alkenyl" refers to a straight hydrocarbon moiety of two to six carbon atoms or a branched hydrocarbon moiety of three to six carbon atoms having at least one carbon-carbon double bond, such as ethenyl (i.e. -CH=CH₂), propenyl, 1-butenyl, 2-butenyl, isobutenyl, 1-pentenyl, 2-methyl-1-butenyl, 1-hexenyl and the like; the term "substituted alkenyl" refers to an alkenyl moiety which is further substituted at a substitutable carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "alkynyl" refers to a straight hydrocarbon moiety of two to six carbon atoms or a branched hydrocarbon moiety of four to six carbon atoms having one carbon to carbon triple bond such as ethynyl

-7-

(i.e. $-C\equiv CH$), 1-propynyl, 1-butynyl, 1-pentynyl, 2-pentynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl and the like; the term "substituted alkynyl" refer to an alkynyl moiety which is further substituted at a substitutable carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "alkoxy" refers to an alkyl moiety containing from 1 to 6 carbon atoms covalently bonded to an adjacent structural element through an oxygen atom, such as for example, methoxy (i.e. $-OCH_3$), ethoxy, propoxy, isopropoxy, butoxy, pentoxy, hexoxy and the like; the term "substituted alkoxy" refers to an alkoxy moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "alkenylalkyl" refers to an alkenyl moiety of two to six carbon atoms covalently bonded to an alkyl moiety of 1 to 6 carbon atoms; the term "substituted alkenylalkyl" refers to an alkenylalkyl moiety which is further substituted at a carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "alkynylalkyl" refers to an alkynyl moiety of two to six carbon atoms covalently bonded to an alkyl moiety of 1 to 6 carbon atoms; the term "substituted alkynylalkyl" refers to an alkynylalkyl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "alkoxyalkyl" refers to an alkoxy moiety of 1 to 6 carbon atoms covalently bonded to an

-8-

alkyl moiety of 1 to 6 carbon atoms; the term "substituted alkoxyalkyl" refers to an alkoxyalkyl moiety which is further substituted at a carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "amino" refers to a primary ($-NH_2$), a secondary or a tertiary amine wherein each hydrogen can be substituted by an alkyl moiety of one to six carbon atoms or by an aryl moiety of six to fifteen carbon atoms.

The term "dialkylaminoalkyl" refers to a nitrogen atom covalently bonded to three alkyl moieties having 1 to 6 carbon atoms in each alkyl moiety, and one alkyl moiety is bonded to an adjacent structural element.

The term "aryl" refers to a carbocyclic moiety containing at least one benzenoid-type ring, with the aryl groups preferably containing from 6 to 15 carbon atoms, for example, phenyl, naphthyl, indenyl, indanyl, and the like; the term "substituted aryl" refers to an aryl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "arylalkyl" refers to an aryl moiety of 6 to 15 carbon atoms covalently bonded to an alkyl moiety of one to six carbon atoms such as, for example, benzyl, phenylethyl, and the like; the term "substituted aralkyl" refers to an aralkyl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "arylalkenyl" refers to an aryl moiety of six to fifteen carbon atoms covalently bonded to an

-9-

alkenyl moiety of two to six carbon atoms, such as, for example, 2-phenyl-1-ethenyl (cinnamyl), 4-phenyl-2-butenyl and the like; the term "substituted arylalkenyl" refers to an arylalkenyl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "arylalkoxy" refers to an aryl moiety of one to six carbon atoms covalently bonded to an alkoxy moiety of one to six carbon atoms; the term "substituted arylalkoxy" refers to an arylalkoxy moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "aryloxy" refers to an aryl moiety covalently bonded through an oxygen (i.e. -O-) atom, such as, for example, phenoxy and the like; the term "substituted aryloxy" refers to an aryloxy moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "alkoxyaryl" refers to an alkoxy moiety of one to six carbon atoms covalently bonded to an aryl moiety of six to 15 carbon atoms, such as, for example 2-methoxyphenyl, 2- or 4-ethoxynaphthyl, 6-propoxyindenyl and the like; the term "substituted alkoxyaryl" refers to an alkoxyaryl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl, or salts thereof, nitro and hydroxyl;

-10-

The term "aryloxyaryl" refers to an aryloxy moiety as defined hereinbefore covalently bonded to an aryl moiety of six to 15 carbon atoms, such as, for example, phenoxyphenyl, 1-naphthyloxyphenyl and the like; the term "substituted aryloxyaryl" refers to an aryloxyaryl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "aryloxyalkyl" refers to an aryloxy moiety as defined hereinbefore covalently bonded to an alkyl moiety of one to six carbon atoms, such as, for example, phenoxymethyl, 1-naphthyloxyethyl and the like; the term "substituted aralkyl" refers to an aralkyl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "acyl" refers to a carbonyl moiety

($\overset{\text{O}}{\parallel}\text{C}-$) bonded to a hydrogen, alkyl, aryl, alkoxy, amino or

an aryloxy group such as a formyl moiety ($\text{H}-\overset{\text{O}}{\parallel}\text{C}-$), an

alkanoyl moiety ($\text{alkyl}-\overset{\text{O}}{\parallel}\text{C}-$) of one to six carbon atoms in

the alkyl portion, an aroyl moiety ($\text{aryl}-\overset{\text{O}}{\parallel}\text{C}-$) of six to 15 carbon atoms in the aryl portion, an ester moiety

($\text{alkoxy}-\overset{\text{O}}{\parallel}\text{C}-$) of one to six carbon atoms in the alkoxy

portion, an amide moiety ($\text{amino}-\overset{\text{O}}{\parallel}\text{C}-$), or an aryloxy moiety. Typical acyl groups include acetyl, benzoyl, ethoxycarbonyl and the like; the term "substituted acyl"

-11-

refers to the alkyl, aryl, alkoxy, amino, aryloxy, alkanoyl, aroyl, ester, amide or aryloxy portion of the acyl moiety which is further substituted at a carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;


The term "aromatic heterocyclic" refers to a cyclic moiety having at least one O, S and/or N heteroatom interrupting the ring structure and having a sufficient number of unsaturated carbon to carbon double bonds, nitrogen to carbon double bonds, and the like, to provide aromatic character, with the aromatic heterocyclic groups preferably containing from 2 to 14 carbon atoms, for example, 2-, 3- or 4-pyridyl, 2- or 3-furyl, 2- or 3-thienyl, 2-, 4- or 5-thiazolyl, 2-, 4- or 5-imidazolyl, 2-, 4- or 5-pyrimidinyl, 2-pyrazinyl, 3- or 4-pyridazinyl, 3-, 5- or 6-[1,2,4-triazinyl], 3- or 5-[1,2,4-thiadiazolyl], 2-, 3-, 4-, 5-, 6- or 7-benzofuranyl, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, and the like; the term "substituted aromatic heterocyclic" refers to an aromatic heterocyclic moiety which is further substituted at a carbon or heteroatom by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "heterocyclic alkyl" refers to an aromatic heterocyclic moiety of 2 to 14 carbon atoms as defined hereinbefore, covalently bonded to an alkyl moiety of one to six carbon atoms; the term "substituted heterocyclic alkyl" refers to a heterocyclic alkyl moiety which is further substituted at a carbon or heteroatom by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano or salts thereof, nitro and hydroxyl;

-12-


The term "heterocyclic cycloalkyl" refers to a saturated carbocyclic ring of two to six carbon atoms having at least one oxygen, sulfur or nitrogen atom, or combination thereof, interrupting the ring structure, and are substituted at a carbon atom, such as 2-, 3-, or 4-piperidyl, 2-dioxanyl, 2- or 3-oxazetidinyl, 2-oxiranyl, or 3-, 4-, 5- or 6-thiazinyl and the like; the term "substituted heterocyclic cycloalkyl" refers to an heterocyclic cycloalkyl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;


The term "heterocyclic cycloalkylalkyl" refers to a heterocyclic cycloalkyl moiety of two to six carbon atoms covalently bonded to an alkyl moiety of one to six carbon atoms; the term "substituted cycloalkalkyl" refers to a cycloalkalkyl moiety which is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "sulfoxide" refers to a sulfoxide moiety (i.e. R-SO-R-) wherein each R independently represents an alkyl moiety of one to six carbon atoms or an aryl moiety of 6 to 15 carbon atoms, such as, for example, $\text{CH}_3\text{-SO-}$ ; the term "substituted sulfoxide" refers to a sulfoxide moiety as defined above where "R" group is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "sulfonyl" refers to a sulfonyl moiety (i.e. R-SO₂-R-) wherein each R independently represents alkyl of one to six carbon atoms or aryl of six to twelve


-13-

carbon atoms, such as, for example, $\text{CH}_3\text{-SO}_2\text{---}$  ; the term "substituted sulfonyl" refers to a sulfonyl moiety as defined above whose "R" group is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "sulfide" refers to a sulfide moiety (i.e. R-S-R-) wherein each R independently represents alkyl of one to six carbon atoms or aryl of six to twelve carbon atoms, such as, for example, alkylthioalkyl, or alkylthioaryl such as $\text{CH}_3\text{-S-}$  ; the term "substituted sulfide" refers to a sulfide moiety as defined above whose "R" group is further substituted at the carbon by one or more of the following groups: halo, alkyl of one to six carbon atoms, aryl of six to fifteen carbon atoms, cyano, carboxyl or salts thereof, nitro and hydroxyl;

The term "hydroxyalkyl" refers to an alkyl moiety in which one or more of the hydrogens is replaced by a hydroxy moiety, such as, for example, hydroxymethyl (i.e. $\text{-CH}_2\text{OH}$), hydroxyethyl, 1-hydroxypropyl, 2-hydroxypropyl, 3-hydroxybutyl, 4-hydroxyhexyl and the like.

The base employed in the present process is any substance which will remove a proton from the hydroxyl (-OH) of the moiety



in order to cyclize the amino alcohol (IX) with the cyano compound (VIII) to give the desired oxazoline compound (X). The base is neither the amino alcohol (IX) nor the cyano compound (VIII).

-14-

Bases which can be employed in process of the present invention can be a non-aqueous base such as lithium diisopropyl amide, lithium hexamethylsilazide, sodium hexamethylsilazide and potassium hexamethylsilazide; or potassium t-butoxide and sodium methoxide. The base can be an alkali metal carbonate such as sodium, potassium, lithium or cesium carbonate or an alkaline earth metal carbonate such as calcium or barium carbonate; hydroxides such as sodium and potassium hydroxides; and hydrides such as sodium or potassium hydrides. The base can also be ammonia (NH_3) or an organic base including urea; a secondary amine such as dimethylamine, diphenylamine, N-methyl N-propylamine, diethylamine, diisopropylamine, N-methylaniline, piperazine, piperidine, pyrrolidine; or a tertiary amine such as trimethylamine, dimethylaniline, N,N-dimethylpropylamine, N,N-dimethylpiperidine, N,N-dimethylbutylamine, triethylamine. The base can also be an heterocyclic nitrogen containing compound such as isoquinoline, morpholine, purine, pyridine, pyrazine, pyrimidine, quinoline or polyvinyl pyridine, preferably DBU. Other bases which may be suitably employed in the present process are disclosed in "Modern Synthetic Reactions" by H. House, W.A. Benjamin, Inc., Menlo Park, California, 1972, 856 pages. Where appropriate, mixtures of any of the above bases can be employed.

The base is used in amounts effective to remove the requisite proton from the hydroxy moiety. Such amounts in terms of mole ratios (moles base:mole amino alcohol (IX)) can range from about 1,000 to 0.001:1, preferably from about 10 to 0.01:1, more preferably from about 1 to 0.1:1, most preferably about 0.15:1.

The alcohol solvent employed in the present invention is preferably dihydric (two OH groups-diols) such as C-2 to C-10 glycols and derivatives. Represen-

-15-

tative C-2 to C-10 glycols and derivatives include ethylene glycol, propylene glycol, 1,2-butanediol, 1,4-butanediol, pentanediols and the like, more preferably ethylene glycol.

The alcohol solvent employed in the present invention is also preferably polyhydric (three or more OH groups-polyols). Representative polyhydric alcohols include glycerol (1,2,3-propanetriol), 1,2,4-butanetriol, penta-erythritol and the like, more preferably glycerol. Where suitable, the process of the present invention can employ a mixture of a dihydric and a polyhydric alcohol, most preferably ethylene glycol and glycerol in a volume ratio of 1:2 (ethylene glycol:glycerol).

The alcohol solvent employed in the present process can be used in amounts which can range from an amount sufficient to at least partially solubilize one or both of the reactants and/or the desired product to an amount in excess of either starting reactant. Generally the amount of alcohol solvent can range from about 1 to 5,000 percent or more by weight of an individual reactant, preferably from about 100 to 1,000 percent by weight, most preferably from about 100 to about 300 percent.

In the process of preparing the oxazoline compound of formula (X), the cyano compound (VIII) is contacted with the aminoalcohol (IX) in amounts and under conditions effective to yield the desired oxazoline compound of formula (X). The cyano compound (VIII) is contacted with the aminoalcohol (IX) at temperatures ranging from about -10 to about 200 degrees Centigrade ($^{\circ}\text{C}$), preferably from about 70 to about 150°C , most preferably from about 100 to about 110°C . The contacting is performed at ambient pressures although pressures greater or less than ambient can be employed. The

-16-


contacting of the reactants can be carried out for about 5 minutes to about 72 hours or more until the reaction is substantially completed, preferably from about 1 hour (hr) to about 48 hours. Also preferred is that the reactants are stirred during the contacting procedures. The cyano compound (VIII) can be contacted with the amino alcohol of (IX) in molar ratios ranging from about 100 to 0.1:1; (moles cyano compound (VIII): mole amino alcohol (IX)), preferably from about 10 to 1:1, most preferably from about 2 to 1:1.

The stereochemistry of the oxazoline compounds (X) is preserved with respect to the stereochemistry of the starting materials. For example, when an amino alcohol of S,S' configuration is contacted with benzonitrile, the resultant oxazoline (X) has a S,S' stereoisomeric configuration.

After the reaction is completed, the desired oxazoline compound (X) can be recovered by conventional separatory and recovery methods such as phase separation, distillation or evaporation of any solvents present, crystallization, chromatography, filtration and the like. For example, the reaction mixture can be diluted with water and the oxazoline compound (X) is recovered by filtration.


Preparation of Starting Materials

The cyano compounds (VIII) are known and can be prepared by conventional procedures, such as for example, by dehydration of the corresponding amide with a dehydrating agent such as phosphorous oxychloride. The dehydration reaction is illustrated as follows:



wherein R^5 is as defined hereinbefore. Similarly, other methods for preparing the cyano compounds (VIII) are disclosed in references such as H.O. House, Modern Synthetic Reactions, Second Edition, W.A. Benjamin Inc., (1979) pp. 79 and 623-628, whose preparation teachings are incorporated herein by reference.

The aminoalcohol compounds (IX) are known and can be prepared by conventional procedures, such as, for example by epoxidation of the corresponding olefin (V) by an epoxidizing method, followed by cleavage of the epoxide ring (VI) by azide to give the azido compound (VII), followed by reduction of the azido compound with a reducing agent to give the requisite aminoalcohol (IX).



-18-

wherein R^1 , R^2 , R^3 and R^4 are as defined hereinbefore. Representative epoxidizing methods include any suitable peracid reactant such as, for example, pertrifluoroacetic acid (CF_3COOH), perbenzoic acid (C_6H_5COOH), peracetic acid (CH_3COOH) and the like, as well as formation of a halohydrin followed by base.

In situations where any of R^1 , R^2 , R^3 or R^4 of olefin compound (V) contains an oxidizable functionality, such as an unsaturated bond, the olefin compound (V) can be selectively epoxidized and/or the requisite epoxide ring compound (VI) thus prepared can be further isolated in order to prepare the azido compound (VII). Representative azides include those of alkali earth metals, such as sodium azide, potassium azide, lithium azide, and the like.

The term "reducing agent" refers to any substance which will furnish electrons by its capacity to lose electrons easily, in order to cause the azido compound (VII) receiving the electrons to be reduced to the desired amino alcohol (IX). Examples of reducing agents include, but are not limited to hydrogenating agents and to metal hydrides, such as lithium aluminum hydride ($LiAlH_4$).

The term "hydrogenating agent" is intended to include the requisite hydrogenating catalyst(s) and hydrogen (H_2) source for reducing the azido compound (VII) to the aminoalcohol (IX). Various selected catalysts and conditions are described in "Catalytic Hydrogenation in Organic Synthesis", (1978) Morris Freifelder, Chapter 4, Olefins, pg. 15-25, John Wiley and Sons. For example, the hydrogenating catalyst can be nickel, palladium, platinum, platinum oxide, platinum on carbon, and mixture thereof.

-19-

Similarly, in situations where R^1 , R^2 , R^3 or R^4 of azido compound (VII) contains a reducible functionality such as an unsaturated bond or a sulfur atom, azido compound (VII) can be reduced with a selective reducing agent and/or the requisite amino alcohol (IX) thus prepared can be further isolated in order to prepare oxazoline compound (X).


Other representative methods of preparing the aminoalcohols of formula (IX) are described in Leroy G. Wade, Jr., Vol. 5, John Wiley and Sons, (1984) pp. 430-431 and in Calvin Buchler and Donald Pearson, Survey of Organic Synthesis, Vol. 1, Wiley Interscience, New York (1970), pp. 226, 466 and 475. The preparative teachings of these references are incorporated herein by reference.

Salts of the aminoalcohol (IX) can be prepared by contacting the aminoalcohol (IX) with organic or inorganic acids. Representative organic acids include but are not limited to oxalic, tartaric, acetic, trifluoroacetic, citric, maleic and the like. Representative inorganic acids include hydrochloric, sulfuric or phosphoric acids in about equimolar amounts or in amounts less than equimolar of the acid relative to the aminoalcohol.

The following examples illustrate the present invention in a manner of which it can be practiced but, as such, should not be construed as limitations upon the overall scope of the same.

SUBSTITUTE SHEET

-20-


EXAMPLE 1PREPARATION OF D-2-PHENYL-4-(4-METHYLMERCAPTOPHENYL)-2-OXAZOLO-5-METHANOL

A suspension of 5 grams (g) (23.5 millimole (mmole)) of D(-)-2-amino-1-(4-methylmercaptophenyl)-1,3-propanediol (available from Fuji Chemical Co., Japan) and 0.5 g (3.6 mmole) of potassium carbonate (K₂CO₃) in 7.5 milliliters (mL) of ethylene glycol and 4.1 mL of glycerol is heated to a temperature of 105°C with stirring. Benzonitrile (4 mL, 39.2 mmole, 1.67 equivalents) is added and the mixture is stirred at 105°C for 18 hr under a blanket of nitrogen gas. Completeness of reaction is determined by thin layer chromatography (TLC) on silica gel (9:1 - methylene chloride: methanol). The product is isolated by precipitation into water, to give 6.8g (purity 97%, 22.0 mmole) a 93 percent yield.

-21-

EXAMPLE 2

PREPARATION OF 5-METHYL-2-PHENYL-2-OXAZOLINE




To a mixture of five mL of 2-aminopropanol (62.8 mmole) and 273 milligrams (mg) of potassium carbonate (37.7 mmole) in ethylene glycol (20.5 mL) and glycerol (11.5 mL) heated to a temperature of 105 degrees °C, is added benzonitrile (9.8 mL; 96.4 mmol)). The reaction mixture is stirred at 105°C for 18 hours, diluted with hexane and washed with water. The solvent and excess benzonitrile from the organic layer are removed by distillation to give 10.45g (purity 85 percent (%)), of title compound, a yield of 92 percent.

-22-

EXAMPLE 3

PREPARATION OF D-2-(p-NITROPHENYL),
4-(4-METHYLMERCAPTOPHENYL),
-2-OXAZOLO-5-METHANOL



To a mixture of 1 g of D(-)-2-amino-1-(4-methylmercaptophenyl)-1,3-propanediol (4.7 mmole) and 0.1g of potassium carbonate in 1.5 mL of ethylene glycol and 0.8 mL of glycerol, heated to a temperature of 100°C, 1.0 g of benzonitrile is added, and stirred for 3 hours. Analysis of the reaction mixture by high pressure liquid chromatography indicates a yield of title compound to be 85 percent.

-23-


While the present invention has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications, and variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

SUBSTITUTE SHEET

-24-

WHAT IS CLAIMED IS:


1. A process for preparing an oxazoline compound of the formula:



comprising contacting a cyano compound of the formula



with an amino alcohol compound of the formula



or salt thereof,




wherein R^1 , R^2 , R^3 , R^4 and R^5 can be the same or different, each independently represents

hydrogen, alkyl, haloalkyl, cycloalkyl or substituted cycloalkyl, cycloalkalkyl or substituted cycloalkalkyl, alkenyl or substituted alkenyl, alkynyl, alkenylalkyl or substituted alkenylalkyl, alkynylalkyl, alkoxyalkyl or substituted alkoxyalkyl, dialkylamino-alkyl, aryl or substituted aryl, arylalkyl or substituted

-25-

arylalkyl, arylalkenyl or substituted arylalkenyl, alkoxyaryl or substituted alkoxyaryl, aryloxyaryl or substituted aryloxyaryl, aryloxyalkyl or substituted aryloxyalkyl, acyl or substituted acyl, aromatic heterocyclic or substituted aromatic heterocyclic, heterocyclic alkyl or substituted heterocyclic alkyl, heterocyclic cycloalkyl or substituted heterocyclic cycloalkyl, heterocyclic cycloalkylalkyl or substituted heterocyclic cycloalkylalkyl, sulfoxide or substituted sulfoxide, sulfonyl or substituted sulfonyl, sulfide or substituted sulfide or hydroxyalkyl, in the presence of a base and a dihydric alcohol solvent, a polyhydric alcohol solvent or mixtures thereof.

2. The process of claim 1 wherein as to the amino alcohol (IX), R^1 and R^2 independently represent hydrogen, sulfide, sulfoxide, sulfonyl, aryl or substituted aryl; and R^3 and R^4 independently represent hydrogen or $-CH_2OH$.

3. The process of claims 1 or 2 wherein R^1 and R^2 independently represent hydrogen or CH_3-S- , CH_3-SO- , CH_3-SO_2- ; and R^3 and R^4 independently represent hydrogen or $-CH_2OH$.

4. The process of claims 1, 2 or 3 wherein as to the cyano compound (VIII) R^5 is phenyl, 4-nitrophenyl, cinnamyl, 4-methoxycinnamyl or dichloromethyl.

5. The process of claims 1, 2, 3 or 4 wherein the base is diazabicycloundecene or potassium carbonate.

6. The process of claims 1, 2, 3, 4 or 5 wherein the base is employed in an amount ranging from about 10 to 0.01 moles base to one mole amino alcohol.

-26-

7. The process of claims 1, 2, 3, 4, 5 or 6 wherein the alcohol solvent is ethylene glycol.
8. The process of claims 1, 2, 3, 4, 5, 6 or 7 wherein the alcohol solvent is a mixture of ethylene glycol and glycerol.
9. The process of claims 1, 2, 3, 4, 5, 6, 7 or 8 wherein the contacting is performed at temperatures ranging from about 70 to about 150°C.
10. The process of claims 1, 2, 3, 4, 5, 6, 7 or 8 wherein the contacting is performed at temperatures ranging from about 100 to about 110°C.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 89/03827

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC ⁵ : C 07 D 263/08		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System ¹	Classification Symbols	
IPC ⁵	C 07 D 263/00	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ¹⁰	Citation of Document, ¹¹ with Indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	Justus Liebigs Annalen der Chemie, no. 6, June 1974, (Weinheim, DE), H. Witte et al.: "Cyclische Imidsäure- ester aus Nitrilen und Aminoalkoholen", Seiten 996-1009 --	1
A	US, A, 2402198 (DONALD J. LODER) 18 June 1946 --	1
A	US, A, 3979405 (ISTVAN TIBOR TOTH) 7 September 1976 --	1
A	EP, A, 0123123 (HENKEL KGaA) 31 October 1984 see claims -----	1
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
20th November 1989	15. 12. 89	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	T.K. WILLIS	

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

US 8903827

SA 31153

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on 06/12/89. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 2402198		None	
US-A- 3979405	07-09-76	AT-B- 331791	25-08-76
		BE-A- 806342	15-02-74
		CA-A- 1021343	22-11-77
		CH-A- 602677	31-07-78
		DE-A- 2352055	09-05-74
		FR-A, B 2203633	17-05-74
		GB-A- 1426028	25-02-76
		NL-A- 7314414	23-04-74
		SE-B- 386173	02-08-76
		US-A- 4119633	10-10-78
EP-A- 0123123	31-10-84	DE-A- 3310905	27-09-84